

SYMMETRY RESULTS FOR CLASSICAL SOLUTIONS OF MONGE-AMPERE SYSTEM IN THE PLANE

LI MA, BAIYU LIU

Dedicated to Louis Nirenberg on the occasion of his 85th birthday

ABSTRACT. In this paper, by the method of moving planes, we prove the symmetry result which says that classical solutions of Monge-Ampere system in the whole plane are symmetric about some point. Our system under consideration comes from the differential geometry problem.

Keyword: Moving plane, positive solutions, radial symmetric, Monge-Ampere system

Mathematics Subject Classification: 35J60, 53C21, 58J05

1. INTRODUCTION

It is an interesting question to consider the relation between two convex surfaces in space. There are at least two famous theorems for convex surfaces in 3-space. One is the Cohn-Vossen theorem, which says that if two closed convex surfaces differ by an isometry, then they are the same by a translation. The other is the Minkowski theorem, which says that if two closed convex surfaces share the same Gauss curvature, then they are differ only by a translation. For a simple and beautiful proof of the latter, one may see the paper of S.S.Chern [4]. We study the relation between two non-compact convex surfaces in space, which leads us to consider a symmetry result for the Monge-Ampere system in the plane $\mathbf{R}^2$. Assume that we are given two functions $K_1(x, u, v)$ and $K_2(x, u, v)$ for $(x, u, v) \in \mathbf{R}^2 \times \mathbf{R}^2$. We are looking for a pair of function $u = u(x)$ and $v = v(x)$ with their graphs $\Gamma(x, u(x))$ and $\Gamma(x, v(x))$ such that $K_1(x, u(x), v(x))$ and $K_2(x, u(x), v(x))$ are Gaussian curvatures of the graphs $\Gamma(x, u(x))$ and $\Gamma(x, v(x))$ respectively. Then we are lead to solve the Monge-Ampere system

$$K_1(x, u, v) = \frac{\det D^2 u(x)}{(1 + |Du(x)|^2)^2}, \quad \text{in } \mathbf{R}^2$$

and

$$K_2(x, u, v) = \frac{\det D^2 v(x)}{(1 + |Dv(x)|^2)^2}, \quad \text{in } \mathbf{R}^2.$$

* The research is partially supported by the National Natural Science Foundation of China 10631020 and SRFDP 20060003002.

Single Monge-Ampere equation (with Dirichlet boundary condition) has been studied by many authors, one may see [3] and [11] for more references.

We study here a little more generalized version of the Monege-Ampere system above. We study the symmetry result for classical solutions of the Monege-Ampere system:

$$(1) \quad \begin{cases} \det(D^2u) + g(u, v, \nabla u) = 0, & \text{in } \mathbf{R}^2, \\ \det(D^2v) + f(u, v, \nabla v) = 0, & \text{in } \mathbf{R}^2, \\ (D^2u) > 0, (D^2v) > 0, & \text{in } \mathbf{R}^2, \\ \lim_{|x| \rightarrow \infty; x_1 < \lambda} (u(x) - u(x^\lambda)) \geq 0, & \text{for } \lambda < 0, \\ \lim_{|x| \rightarrow \infty; x_1 < \lambda} (v(x) - v(x^\lambda)) \geq 0, & \text{for } \lambda < 0, \\ \lim_{|x| \rightarrow \infty; x_1 > \lambda} (u(x) - u(x^\lambda)) \geq 0, & \text{for } \lambda > 0, \\ \lim_{|x| \rightarrow \infty; x_1 > \lambda} (v(x) - v(x^\lambda)) \geq 0, & \text{for } \lambda > 0, \\ \lim_{|x| \rightarrow \infty} u(x) = \lim_{|x| \rightarrow \infty} v(x) = +\infty, \\ \lim_{|x| \rightarrow \infty} |\nabla u(x)| = \lim_{|x| \rightarrow \infty} |\nabla v(x)| = \infty, \end{cases}$$

for $x^\lambda = (2\lambda - x_1, x_2)$ is the reflection of the point x with respect to any line $T_\lambda = \{x_1 = \lambda\}$, where $f, g \in C^1(\mathbf{R}^2 \times \mathbf{R}^2, \mathbf{R})$.

We suppose that:

- (i) f and g are symmetric in $p_1 : \forall (u, v) \in \mathbf{R}^2, \forall p_1, p_2 \in \mathbf{R}$,
 $g(u, v, p_1, p_2) = g(u, v, -p_1, p_2), f(u, v, p_1, p_2) = f(u, v, -p_1, p_2);$
- (ii) $C_1 > \frac{\partial g}{\partial v}(u, v, p_1, p_2) > 0, \quad C_2 > \frac{\partial f}{\partial u}(u, v, p_1, p_2) > 0,$
 $\forall (u, v) \in \mathbf{R}^2, \forall p_1, p_2 \in \mathbf{R};$
- (iii) $\lim_{|u|^{-1} + |v|^{-1} + |(p_1, p_2)|^{-1} \rightarrow 0} \frac{\partial g}{\partial u}(u, v, p_1, p_2) < 0,$
 $\lim_{|u|^{-1} + |v|^{-1} + |(p_1, p_2)|^{-1} \rightarrow 0} \frac{\partial f}{\partial v}(u, v, p_1, p_2) < 0;$
- (iv) $\lim_{|u|^{-1} + |v|^{-1} + |(p_1, p_2)|^{-1} \rightarrow 0} \det \begin{pmatrix} \frac{\partial g}{\partial u} & \frac{\partial g}{\partial v} \\ \frac{\partial f}{\partial u} & \frac{\partial f}{\partial v} \end{pmatrix} (u, v, p_1, p_2) > 0,$

where C_i are positive constants.

Our result is the following.

Theorem 1. *Let (u, v) be a classical solution of system (1). Suppose (i)-(iv) hold, then the solution (u, v) is symmetric in x_1 direction, i.e., there is some λ such that $u(x_1, x_2) = u(2\lambda - x_1, x_2)$ and $v(x_1, x_2) = v(2\lambda - x_1, x_2)$.*

Notation: We shall use $(\overline{A, B})$ to denote the open interval in the line from A to B .

The method to prove Theorem 1 is motivated by [2], [8], [10], and [9]. We use the method of moving planes, which has been used in [5], [1], [6], and [7].

2. MONGE-AMPERE SYSTEM IN $\mathbf{R}^2$

We prove our theorem in this section.

In what follows, we shall use the method of moving plane. To proceed, we start by considering lines parallel to $x_1 = 0$, coming from $-\infty$. For each $\lambda \in \mathbf{R}$, we define

$$\Sigma_\lambda := \{x \in \mathbf{R}^2 \mid x_1 < \lambda\}, \quad T_\lambda := \partial\Sigma_\lambda = \{x \in \mathbf{R}^2 \mid x_1 = \lambda\}.$$

For any point $x = (x_1, x_2) \in \Sigma_\lambda$, let $x^\lambda = (2\lambda - x_1, x_2)$ be the reflected point with respect to the line T_λ . We define the reflected functions by

$$u_\lambda(x) := u(x^\lambda), \quad v_\lambda(x) := v(x^\lambda),$$

and introduce the functions

$$U_\lambda(x) := u_\lambda(x) - u(x), \quad V_\lambda(x) := v_\lambda(x) - v(x).$$

By the definition of u_λ , $\det(D^2 u_\lambda)(x) = \det(D^2 u)(x^\lambda)$. By using the integral form of the theorem of the mean, we obtain:

$$(2) \quad \det(D^2 u_\lambda) - \det(D^2 u) = a_{ij} U_{\lambda,ij},$$

where

$$(3) \quad (a_{ij}(x)) = \frac{1}{2}(\det(D^2 u_\lambda)(D^2 u_\lambda)^{-1} + \det(D^2 u)(D^2 u)^{-1}).$$

Noticing that the solution (u, v) are convex everywhere, we have

$$(4) \quad (a_{ij}(x)) > 0, \quad x \in \Sigma_\lambda.$$

On the other hand, by using (i), we have

$$\begin{aligned} (5) \quad \det(D^2 u_\lambda)(x) &= -g(u, v, \nabla u)(x^\lambda) \\ &= -g(u_\lambda, v_\lambda, -u_{\lambda,1}, u_{\lambda,2}) \\ &= -g(u_\lambda, v_\lambda, \nabla u_\lambda). \end{aligned}$$

Similarly we have the elliptic equation for V_λ with the elliptic coefficient matrix $(b_{ij}) = (b_{ij}(x))$.

Therefore, we find that (U_λ, V_λ) satisfies an elliptic linear system: in Σ_λ

$$\begin{aligned} (6) \quad a_{ij} U_{\lambda,ij} &+ \frac{\partial g}{\partial p_1}(u_\lambda, v_\lambda, \theta_1(x, \lambda), u_{\lambda,2}) U_{\lambda,1} + \frac{\partial g}{\partial p_2}(u_\lambda, v_\lambda, u_{\lambda,1}, \theta_2(x, \lambda)) U_{\lambda,2} \\ &+ \frac{\partial g}{\partial u}(\xi_1(x, \lambda), v, \nabla u) U_\lambda + \frac{\partial g}{\partial v}(u_\lambda, \eta_1(x, \lambda), \nabla u) V_\lambda = 0, \end{aligned}$$

(7)

$$b_{ij}V_{\lambda,ij} + \frac{\partial f}{\partial p_1}(u_\lambda, v_\lambda, \tau_1(x, \lambda), v_{\lambda,2})V_{\lambda,1} + \frac{\partial f}{\partial p_2}(u_\lambda, v_\lambda, v_{\lambda,1}, \tau_2(x, \lambda))V_{\lambda,2} \\ + \frac{\partial f}{\partial u}(\xi_2(x, \lambda), v_\lambda, \nabla v)U_\lambda + \frac{\partial f}{\partial v}(u, \eta_2(x, \lambda), \nabla v)V_\lambda = 0,$$

where for $i = 1, 2$,

$$(8) \quad \xi_i(x, \lambda) \in \overline{(u(x), u_\lambda(x))}, \quad \eta_i(x, \lambda) \in \overline{(v(x), v_\lambda(x))},$$

$$(9) \quad \theta_i(x, \lambda) \in \overline{(u_i(x), u_{\lambda,i}(x))},$$

$$(10) \quad \tau_i(x, \lambda) \in \overline{(v_i(x), v_{\lambda,i}(x))}.$$

We define

$$\Sigma_\lambda^{U^-} := \{x \in \Sigma_\lambda \mid U_\lambda(x) > 0\}, \quad \Sigma_\lambda^{V^-} := \{x \in \Sigma_\lambda \mid V_\lambda(x) > 0\}.$$

For simplicity, we first give two lemmata.

Lemma 2. *Assume (i)-(iv) hold. Then there exists a constant $R_1 > 0$, such that*

$$\forall x \in \{\mathbf{R}^2 \setminus B_{R_1}(0)\} \cap \Sigma_\lambda^{U^-}; \quad \forall y \in \{\mathbf{R}^2 \setminus B_{R_1}(0)\} \cap \Sigma_\lambda^{V^-}$$

the following hold

$$\frac{\partial g}{\partial u}(z, v(x), \nabla u(x)) < 0, \quad \frac{\partial f}{\partial v}(u(y), w, \nabla v(y)) < 0,$$

where $z \in \overline{(u_\lambda(x), u(x))}$ and $w \in \overline{(v_\lambda(y), v(y))}$ arbitrarily.

Proof. Using (iii), we choose $\epsilon > 0$ such that $\frac{\partial g}{\partial u}(u, v, p_1, p_2) < 0$, if $|u|^{-1} + |v|^{-1} + |(p_1, p_2)|^{-1} < \epsilon$. For this particular ϵ , since $\lim_{|x| \rightarrow +\infty} (|u(x)|^{-1} + |\nabla u|^{-1}) = 0$ and $\lim_{|x| \rightarrow +\infty} |v(x)|^{-1} = 0$, there exists a constant $R_1 > 0$, such that $|u(x)|^{-1} + |v(x)|^{-1} + |\nabla u|^{-1}(x) < \epsilon$, if $|x| > R_1$. For all $x \in \{\mathbf{R}^2 \setminus B_{R_1}(0)\} \cap \Sigma_\lambda^{U^-}$, $|u(x)|^{-1} + |v(x)|^{-1} + |\nabla u|^{-1}(x) < \epsilon$. Notice that in this case $z \in \overline{(u_\lambda(x), u(x))} = (u(x), u_\lambda(x))$. Thus, $|z|^{-1} + |v(x)|^{-1} + |\nabla u|^{-1}(x) < \epsilon$, $\forall z \in \overline{(u_\lambda(x), u(x))}$. Therefore, $\frac{\partial g}{\partial u}(z, v(x), \nabla u(x)) < 0$, $\forall z \in \overline{(u_\lambda(x), u(x))}$. Similarly, when R_1 is sufficiently large, $\frac{\partial f}{\partial v}(u(y), w, \nabla v(y)) < 0$, if $y \in \{\mathbf{R}^2 \setminus B_{R_1}(0)\} \cap \Sigma_\lambda^{V^-}$. $\square$

Lemma 3. *Assume (i)-(iv) hold. Then there exists a constant $R_2 > 0$, such that*

$$\forall y_1, y_2 \in \{\mathbf{R}^2 \setminus B_{R_2}(0)\} \cap \Sigma_\lambda^{U^-} \cap \Sigma_\lambda^{V^-},$$

the following holds:

$$(11) \quad \begin{aligned} & \frac{\partial g}{\partial u}(z_1, v(y_1), \nabla u(y_1)) \frac{\partial f}{\partial v}(u(y_2), w_2, \nabla v(y_2)) \\ & - \frac{\partial g}{\partial v}(u_\lambda(y_1), w_1, \nabla u(y_1)) \frac{\partial f}{\partial u}(z_2, v_\lambda(y_2), \nabla v(y_2)) > 0, \end{aligned}$$

where $z_i \in \overline{(u_\lambda(y_i), u(y_i))}$, $w_i \in \overline{(v_\lambda(y_i), v(y_i))}$ ($i = 1, 2$) arbitrarily.

Proof. Since $f, g \in C^1(\mathbf{R}^2 \times \mathbf{R}^2, \mathbf{R})$ and the fact that the solutions growing to infinity at infinity, for all $R \geq 0$, $\forall y_1, y_2 \in \{\mathbf{R}^2 \setminus B_R(0)\} \cap \Sigma_\lambda^{U^-} \cap \Sigma_\lambda^{V^-}$, $\frac{\partial g}{\partial u}(z_1, v(y_1), \nabla u(y_1))$ and $\frac{\partial g}{\partial v}(z_1, v(y_1), \nabla u(y_1))$ are bounded. Using the continuity of $\frac{\partial f}{\partial u}$ and $\frac{\partial f}{\partial v}$, we have

$$\begin{aligned} & \frac{\partial g}{\partial u}(z_1, v(y_1), \nabla u(y_1)) \frac{\partial f}{\partial v}(u(y_2), w_2, \nabla v(y_2)) \\ & - \frac{\partial g}{\partial v}(u_\lambda(y_1), w_1, \nabla u(y_1)) \frac{\partial f}{\partial u}(z_2, v_\lambda(y_2), \nabla v(y_2)) \\ & \geq \frac{1}{2} \left\{ \frac{\partial g}{\partial u}(z_1, v(y_1), \nabla u(y_1)) \frac{\partial f}{\partial v}(u(y_1), v(y_1), \nabla v(y_1)) \right. \\ & \quad \left. - \frac{\partial g}{\partial v}(u(y_1), v(y_1), \nabla u(y_1)) \frac{\partial f}{\partial u}(u(y_1), v(y_1), \nabla v(y_1)) \right\} \\ & \rightarrow \lim_{|u|^{-1} + |v|^{-1} + |(p_1, p_2)|^{-1} \rightarrow 0} \frac{1}{2} \left(\frac{\partial g}{\partial u} \frac{\partial f}{\partial v} - \frac{\partial g}{\partial v} \frac{\partial f}{\partial u} \right)_{(u, v, p_1, p_2)} > 0 \quad (\text{as } R \rightarrow +\infty), \end{aligned}$$

which proves the lemma. $\square$

We now give our Theorem 1, which will be proved by the use of the maximum principle and the method of moving planes as in [2].

Proof. We just have to show that for $\lambda = 0$ there hold

$$U_\lambda(x) \equiv V_\lambda(x) \equiv 0, \quad \forall x \in \Sigma_\lambda.$$

In order to show this, we will apply the moving planes methods in three steps.

Step 1: There exists $\lambda^* < 0$ such that $U_\lambda \leq 0$ and $V_\lambda \leq 0$ in Σ_λ , for all $\lambda \leq \lambda^*$.

Fix $\lambda^* < -\max\{R_1, R_2\}$, then for all $\lambda < \lambda^*$, we have $\forall x \in \Sigma_\lambda$, $|x| > \max\{R_1, R_2\}$. Assume for contradiction that there is a $\lambda < \lambda^*$ and a point $y_0 \in \Sigma_\lambda$, such that $U_\lambda(y_0) > 0$.

Since $\lim_{|x| \rightarrow +\infty} U_\lambda(x) = 0$ and $U_\lambda(x) \equiv 0$, $x \in T_\lambda = \partial\Sigma_\lambda$, we may take $y_1 \in \Sigma_\lambda$, such that

$$U_\lambda(y_1) = \max_{y \in \Sigma_\lambda} U_\lambda(y) > 0.$$

At point y_1 , we have $(U_{\lambda,ij})(y_1) \leq 0$ and $\nabla U_\lambda(y_1) = 0$. Thus, (6) turns to be

$$(12) \quad \frac{\partial g}{\partial u}(\xi_1(y_1, \lambda), v(y_1), \nabla u(y_1))U_\lambda(y_1) + \frac{\partial g}{\partial v}(u_\lambda(y_1), \eta_1(y_1, \lambda), \nabla u(y_1))V_\lambda(y_1) \geq 0.$$

Notice that $y_1 \in \{\mathbf{R}^2 \setminus B_{R_1}(0)\} \cap \Sigma_\lambda^{U^-}$, by using Lemma 2, we have

$$\frac{\partial g}{\partial v}(u_\lambda(y_1), \eta_1(y_1, \lambda), \nabla u(y_1))V_\lambda(y_1) > 0.$$

Combining this with the assumption (ii), we have $V_\lambda(y_1) > 0$, i.e. $y_1 \in \{\mathbf{R}^2 \setminus B_{R_2}(0)\} \cap \Sigma_\lambda^{U^-} \cap \Sigma_\lambda^{V^-}$. On the other hand, we also have

$$\lim_{|x| \rightarrow +\infty} V_\lambda(x) = 0 \quad \text{and} \quad V_\lambda(x) = 0 \quad \forall x \in T_\lambda,$$

hence, we can take $y_2 \in \Sigma_\lambda$ such that

$$V_\lambda(y_2) = \max_{y \in \Sigma_\lambda} V_\lambda(y) > 0.$$

We can repeat the above argument for (7) and show that $U_\lambda(y_2) > 0$ i.e. $y_2 \in \{\mathbf{R}^2 \setminus B_{R_2}(0)\} \cap \Sigma_\lambda^{U^-} \cap \Sigma_\lambda^{V^-}$ and

$$(13) \quad \frac{\partial f}{\partial u}(\xi_2(y_2, \lambda), v_\lambda(y_2), \nabla v(y_2))U_\lambda(y_2) + \frac{\partial f}{\partial v}(u(y_2), \eta_2(y_2, \lambda), \nabla v(y_2))V_\lambda(y_2) \geq 0.$$

Let us put

$$\begin{aligned} J_{11}(\lambda) &= \frac{\partial g}{\partial u}(\xi_1(y_1, \lambda), v(y_1), \nabla u(y_1)) < 0, \\ J_{12}(\lambda) &= \frac{\partial g}{\partial v}(u_\lambda(y_1), \eta_1(y_1, \lambda), \nabla u(y_1)) > 0, \\ J_{21}(\lambda) &= \frac{\partial f}{\partial u}(\xi_2(y_2, \lambda), v_\lambda(y_2), \nabla v(y_2)) > 0, \\ J_{22}(\lambda) &= \frac{\partial f}{\partial v}(u(y_2), \eta_2(y_2, \lambda), \nabla v(y_2)) < 0. \end{aligned}$$

By using (12) and (13), we obtain

$$\begin{aligned} U_\lambda(y_1) &\leq -\frac{J_{12}(\lambda)}{J_{11}(\lambda)}V_\lambda(y_1) \\ &\leq -\frac{J_{12}(\lambda)}{J_{11}(\lambda)}V_\lambda(y_2) \\ &\leq \frac{J_{12}(\lambda)J_{21}(\lambda)}{J_{11}(\lambda)J_{22}(\lambda)}U_\lambda(y_2) \\ &\leq \frac{J_{12}(\lambda)J_{21}(\lambda)}{J_{11}(\lambda)J_{22}(\lambda)}U_\lambda(y_1), \end{aligned}$$

which implies

$$J_{11}(\lambda)J_{22}(\lambda) - J_{12}(\lambda)J_{21}(\lambda) \leq 0.$$

However, as we have shown $y_1, y_2 \in \{\mathbf{R}^2 \setminus B_{R_2}(0)\} \cap \Sigma_\lambda^{U^-} \cap \Sigma_\lambda^{V^-}$, by the result of Lemma 3,

$$J_{11}(\lambda)J_{22}(\lambda) - J_{12}(\lambda)J_{21}(\lambda) > 0,$$

which leads to a contradiction. **Step 1 is completed.**

We now move the plane T_λ toward right, i.e. increase the value of λ , as long as both $U_\lambda(x) \leq 0$ and $V_\lambda(x) \leq 0$ hold. Define:

$$(14) \quad \lambda_0 = \sup\{\lambda \in \mathbf{R} \mid \forall \mu \leq \lambda, U_\mu \leq 0, V_\mu \leq 0, \forall x \in \Sigma_\mu\}.$$

Step 1 implies that $\lambda_0 > -\infty$. On the other hand, as $u(x) > u(0)$, $\forall |x| > R$, for some sufficiently large R , $\lambda_0 < +\infty$. We want to show that $\lambda_0 \geq 0$. Assume not, i.e., $\lambda_0 < 0$.

Since all objects we consider are continuous with respect to λ , we know that

$$U_{\lambda_0} \leq 0 \quad \text{and} \quad V_{\lambda_0} \leq 0, \quad \text{in } \Sigma_{\lambda_0}.$$

Then it follows from (6) (7) and (ii) that in Σ_{λ_0} :

$$(15) \quad \begin{aligned} a_{ij}U_{\lambda_0,ij} &+ \frac{\partial g}{\partial p_1}(u_{\lambda_0}, v_{\lambda_0}, \theta_1(x, \lambda_0), u_{\lambda_0,2})U_{\lambda_0,1} \\ &+ \frac{\partial g}{\partial p_2}(u_{\lambda_0}, v_{\lambda_0}, u_{\lambda_0,1}, \theta_2(x, \lambda_0))U_{\lambda_0,2} + \frac{\partial g}{\partial u}(\xi_1(x, \lambda_0), v, \nabla u)U_{\lambda_0} \geq 0, \end{aligned}$$

$$(16) \quad \begin{aligned} b_{ij}V_{\lambda_0,ij} &+ \frac{\partial f}{\partial p_1}(u_{\lambda_0}, v_{\lambda_0}, \tau_1(x, \lambda_0), v_{\lambda_0,2})V_{\lambda_0,1} \\ &+ \frac{\partial f}{\partial p_2}(u_{\lambda_0}, v_{\lambda_0}, v_{\lambda_0,1}, \tau_2(x, \lambda_0))V_{\lambda_0,2} + \frac{\partial f}{\partial v}(u, \eta_2(x, \lambda_0), \nabla v)V_{\lambda_0} \geq 0, \end{aligned}$$

where θ_i, τ_i, η_i are the same as in (6) and (7).

Step 2. Either $U_{\lambda_0} \equiv 0$ or $V_{\lambda_0} \equiv 0$ hold in Σ_{λ_0} .

Suppose $U_{\lambda_0} \not\equiv 0$ and $V_{\lambda_0} \not\equiv 0$, in Σ_{λ_0} . Thus we can choose $R_3 > \max\{R_1, R_2\}$ s.t. $U_{\lambda_0} \not\equiv 0$ and $V_{\lambda_0} \not\equiv 0$ in $\Sigma_{\lambda_0} \cap B_{R_3}(0)$.

The definition of λ_0 implies that there exist sequences $\{\lambda_k\}_{k=1}^\infty \subset \mathbf{R}$ and $\{y_k\}_{k=1}^\infty \subset \mathbf{R}^2$ such that $\lambda_k > \lambda_0$, $\lim_{k \rightarrow \infty} \lambda_k = \lambda_0$, $y_k \in \Sigma_{\lambda_k}$, and either $U_{\lambda_k}(y_k) > 0$, or $V_{\lambda_k}(y_k) > 0$. Taking $U_{\lambda_k}(y_k) > 0$ (up to a subsequence) as an example, we can rename y_k to be the maximum points, i.e.

$$(17) \quad U_{\lambda_k}(y_k) = \max_{x \in \Sigma_{\lambda_k}} U_{\lambda_k}(x) > 0, \quad k = 1, 2, \dots$$

Then there are two cases could happen.

Case 1. The sequence $\{y_k\}_{k=1}^\infty$ contains a bounded subsequence.

Without loss of generality, we assume $\{y_k\}_{k=1}^\infty \subset \{\Sigma_{\lambda_0} \cap B_{R_3-1}(0)\}$.

$$\lim_{k \rightarrow \infty} y_k = y_0, \quad y_0 \in \cap_{k=1}^{+\infty} \Sigma_{\lambda_k} = \overline{\Sigma_{\lambda_0}}.$$

Noticing that since (a_{ij}) is strictly convex in $\Sigma_{\lambda_0} \cap B_{R_3}(0)$, we can apply the Maximum Principle and the Hopf's lemma to (15) in $\{\Sigma_{\lambda_0} \cap B_{R_3}(0)\}$.

From (17), we have $U_{\lambda_k}(y_k) > 0$, $D^2 U_{\lambda_k}(y_k) \geq 0$, and $\nabla U_{\lambda_k}(y_k) = 0$, which imply

$$(18) \quad U_{\lambda_0}(y_0) \geq 0, \quad D^2 U_{\lambda_0}(y_0) \geq 0, \quad \text{and} \quad \nabla U_{\lambda_0}(y_0) = 0,$$

Since $U_{\lambda_0}(x) \leq 0 \quad \forall x \in \Sigma_{\lambda_0}$, we have $U(y_0) = 0$. Using the Maximum Principle, we have either (a) $U_{\lambda_0} \equiv 0$ in $\{\Sigma_{\lambda_0} \cap B_{R_3}(0)\}$, or (b) $U_{\lambda_0} < 0 \quad \forall x \in \{\Sigma_{\lambda_0} \cap B_{R_3}(0)\}$. Since we haven chosen R_3 to be a large constant such that (a) does not happen, (b) must hold. Therefore, $y_0 \in \partial\{\Sigma_{\lambda_0} \cap B_{R_3}(0)\}$. By using Hopf's lemma, we have $\frac{\partial U_{\lambda_0}}{\partial \nu}(y_0) > 0$, (ν is the outward normal vector on $\partial\{\Sigma_{\lambda_0} \cap B_{R_3}(0)\}$), which contradicts the second equation in (18).

Case 2. $\lim_{k \rightarrow \infty} |y_k| = +\infty$.

In this case, we can choose $k^* > 0$ such that if $k > k^*$, $|y_k| > \max\{R_1, R_2\}$, where R_1, R_2 as in Lemma 2 and Lemma 3. For any fixed $k > k^*$, exactly as in Step 1, we can show $V_{\lambda_k}(y_k) > 0$. After choosing

$$V_{\lambda_k}(x_k) = \max_{y \in \Sigma_{\lambda_k}} V_{\lambda_k}(y) > 0,$$

we also have $U_{\lambda_k}(x_k) > 0$. Therefore,

$$x_k, y_k \in \{\mathbf{R}^2 \setminus B_{R_2}(0)\} \cap \Sigma_{\lambda_k}^{U^-} \cap \Sigma_{\lambda_k}^{V^-}.$$

Similarly defining

$$J_{11}(\lambda_k) = \frac{\partial g}{\partial u}(\xi_1(y_k, \lambda_k), \nu(y_k), \nabla u(y_k)) < 0,$$

$$J_{12}(\lambda_k) = \frac{\partial g}{\partial \nu}(u_{\lambda_k}(y_k), \eta_1(y_k, \lambda_k), \nabla u(y_k)) > 0,$$

$$J_{21}(\lambda_k) = \frac{\partial f}{\partial u}(\xi_2(x_k, \lambda_k), \nu_{\lambda}(x_k), \nabla v(x_k)) > 0,$$

$$J_{22}(\lambda_k) = \frac{\partial f}{\partial \nu}(u(x_k), \eta_2(x_k, \lambda_k), \nabla v(x_k)) < 0,$$

same as in Step 1, we can show $J_{11}(\lambda_k)J_{22}(\lambda_k) - J_{12}(\lambda_k)J_{21}(\lambda_k) \leq 0$, which contradicts Lemma 3. **Step 2 is completed.**

Step 3. Both $U_{\lambda_0} \equiv 0$ and $V_{\lambda_0} \equiv 0$ hold in Σ_{λ_0} .

From Step 2, we let $U_{\lambda_0} \equiv 0$ for example, i.e. $u(x)$ is symmetry about T_{λ_0} . Putting U_{λ_0} into (6) and using the assumption (ii), we obtain $V_{\lambda_0} \equiv 0$. Assume that $\lambda_0 \geq 0$. Doing the same for planes from the right, we know that there is some λ_1 such that both $U_{\lambda_1} \equiv 0$ and $V_{\lambda_1} \equiv 0$ hold in $\Sigma_{\lambda_1}^c$. **Step 3 is completed.**

□

REFERENCES

- [1] H. Berestycki and L. Nirenberg, *On the method of moving planes and the sliding method*, Bol. Soc. Brasil. Mat. (N.S.) 22 (1991), no. 1, 1-37
- [2] J. Busca, B. Sirakov, *Symmetry results for semilinear elliptic systems in the whole space*, J. Differential Equations, 163, 2000, 41-56.
- [3] L. Caffarelli, Yanyan Li, *An extension to a theorem of Jurgens, Calabi, and Pogorelov*, Communications on Pure and Applied Mathematics, Volume 56 Issue 5, Pages 549 - 583
- [4] S.S.Chern, *A proof of the uniqueness of Minkowski problem for convex surfaces*, American journal of mathematics, 79(1957)949-950.
- [5] B. Gidas, W.M. Ni, L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys., 68, (1979), p. 209-243.
- [6] W. Chen, C. Li, *Classification of solutions of some nonlinear elliptic equations*, Duke Math. J., 63(1991) 615-622.
- [7] C. Li, *Monotonicity and symmetry of solutions of fully nonlinear elliptic equations on bounded domains*, Comm. in Partial Differential Equations, 16(1991) 491-529.
- [8] C. Li, *Monotonicity and symmetry of solutions of fully nonlinear elliptic equations on unbounded domains*, Comm. in Partial Differential Equations, 16(1991) 585-615.
- [9] Li Ma, *Three remarks on mean field equations*, to appear in Pacific Journal of Mathematics.
- [10] Li Ma, B.Liu, *Symmetry Results for decay solutions of Elliptic Systems in the Whole Space*, preprint, 2009.
- [11] N.S.Trudinger, J.Urbas, *The Dirichlet problem for the equation of prescribed Gauss curvature*, Bull. Aust.Math. Soc., 28(1983)217-231.

LM: DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA
E-mail address: lma@math.tsinghua.edu.cn

BL: DEPARTMENT OF MATHEMATICAL SCIENCES, TSINGHUA UNIVERSITY, BEIJING 100084, CHINA
E-mail address: liuby05@mails.tsinghua.edu.cn